

[1]



## HIGH RESIDUE CONSERVATION TILLAGE FOR ROW CROPS

The autumn 2005 issue of *Inside CEFS* offered a glimpse of our initial experiences using the mechanical roller-crimper for residue management in cotton and soybean. These trials were initiated in 2003 and continued for two growing seasons. Our objectives were to evaluate weed suppression, residue management, and cotton lint and soybean yield response in a high-residue conservation tillage system to determine (1) the physical effect of surface pressed, intact residue and residue orientation on early-season weed suppression using different weed control programs; (2) the relationship between residue decomposition and incident weed pressure; and (3) the effect of residue management on soybean and cotton stand establishment, growth, and yield. Further testing was done in 2005 to assess the effect of tillage and residue management on early season soil moisture in selected treatments. Readers who are not acquainted with the mechanical roller-crimper or the motives behind this project, may review our [first installment](#) from the conservation tillage 'underground' for details. Our present purpose is to summarize results relevant to the project's objectives, as well as lay the foundation for future work in this relatively new and critical area of conservation tillage.

### Methods

In autumn 2003, a rye (cv. 'Wrens Abruzzi') cover crop was established at two sites with similar weed management histories at the NCDA Cherry Farm Research Unit in Goldsboro, NC. Soils at both sites were nearly level, well-drained and moderately permeable Wickham sandy loam. Prior to rye growth termination, biomass production was estimated using 0.5-m<sup>2</sup> quadrates. Rye growth was terminated mechanically, or with glyphosate (1.75 L ha<sup>-1</sup>), and the standing rye residue flattened using one pass of a mechanical roller-crimper, specially built for this project by the Kelley Manufacturing Company of Tifton, GA. USA.

Full-season soybean (Pioneer 95B97) was planted with a six-row John Deere Maximerge vacuum planter calibrated to deliver 344,300 seeds ha<sup>-1</sup> (8 seeds/ft) on 76-cm (30-inch) rows (12-row plots). Cotton (DP 451) was planted using a four-row planter calibrated at 119,170 seeds ha<sup>-1</sup> (3.5 seeds/ft) on 96-cm (38-inch) rows (8-row plots). Weed management programs were selected to represent a range of herbicide inputs, including: (1) rye residue + no herbicide; (2) rye + glyphosate only for burndown; (3) rye + glyphosate + pre-emergent residual herbicide; (4) rye + glyphosate + pre- + post-emergent over-the-top herbicide. Pre- and post-emergence herbicide treatments for soybean (metolachlor + imazaquin pre-broadcast; bentazon + sethoxydim post-broadcast) and cotton (fluometuron pre-broadcast; MSMA post-directed) were applied at NCDA recommended rates.

A no-tillage planting system was used in all plots but cotton, where a subsoil-strip till treatment was introduced. A conventional 'clean' disk tillage treatment and, a no-tillage treatment with cotton and soybean planted in non-rolled, glyphosate-terminated rye, was included in both trials. All treatments were randomized before planting in complete blocks with four replications.

### Weeds

Weed pressure was assessed at-planting (AP), and at two weeks and four weeks post-emergence (PE) by counting the number and species of germinated weeds within three independent 0.5-m<sup>2</sup> quadrates centered on, and coincident to, the nearest plant row occurring at the terminus of alternating left and right side offsets approximately 2 m perpendicular to a diagonal transect in each plot. Total weed biomass was estimated at eight weeks PE by harvesting the above-ground biomass in the same three 0.5m<sup>2</sup> quadrates noted above. The biomass was oven-dried at 70° C to a constant weight. Weed count and biomass data underwent analysis of variance in the MIXED procedure in SAS. A 0.05 level of probability was adopted for testing hypotheses.

### Rye Residue

Residue decomposition was evaluated by placing folded, intact rye residue in 2-mm nylon mesh bags at rates equivalent to field conditions. The bags were retrieved at 2, 4, 6, 8, and 16 weeks after planting. Per cent original residue remaining in the bags was computed and the data fit to an appropriate model using the NLIN (non-linear) procedure in SAS.

### Soil moisture

Three no-tillage treatments were selected for soil moisture testing: (1) NT—no roll + gly + pre + post herb; (2) NT—roll + gly + pre + post herb; and (3) NT — roll + no herb, and two subsoil strip-tillage treatments (1) SST + roll + pre + post herbicide; and (2) SST + roll + no herbicide. Soil was sampled to 45 cm deep in the undisturbed, untrafficked interrow in all no-tillage and disk tillage plots approximately two weeks after crop emergence using a 1.25 cm diameter

Giddings tube. This procedure was modified for subsoil strip-tillage, where two row positions were sampled (1) tilled strip; and (2) undisturbed, untrafficked interrow. Two cores were obtained at each of three depths from each plot and sectioned into 0-15, 15-30, and 30-45-cm depth increments. Two bulk density cores 7.62 cm high x 7.62 cm diameter, were also obtained at each of the same three depths using a Uhland drive head and hammer assembly. Soil moisture content was determined gravimetrically (g H<sub>2</sub>O/g soil) and converted to a volumetric basis (m<sup>3</sup> H<sub>2</sub>O m<sup>3</sup> soil) using bulk density values.

#### Yield

Cotton and soybean were machine harvested and yield determined. Due to high weed pressure in the non-herbicide treated cotton plots in 2005, these required hand harvesting. In this report, metric and English units are presented for comparison of yield.

count AP in two of four observations over two seasons. To compare the effect of rye residue vs. herbicide on post emergence weed suppression, statistical contrasts were employed (**Table 1**). The contrasts showed that weed count and weed biomass were significantly greater ( $p < .0001$ ) when mechanically rolled rye was substituted for herbicide. Moreover, mean weed density increased significantly over time ( $p < .0001$ ), indicating that surface pressed residue along with minimal soil disturbance at planting did not prevent weeds from germinating. Weed count and weed biomass tended to be slightly greater where strip-tillage was applied without herbicide, compared to the no-herbicide, no-tillage treatment. Not surprisingly, summer annuals accounted for the majority of weed species occurring in both no-tillage and subsoil strip-tillage planting systems. Dominant summer annuals included at least four species of pigweed (*Amaranthus* spp.), three species of morning glory

**Table 1.** Mean weed count and biomass under contrasting weed management and tillage over two cotton (COT) and soybean (SOY) cropping seasons at Goldsboro, NC.

TREATMENT	CONTRAST <sup>‡</sup>	WEED DENSITY, ct m <sup>-2</sup>						WEED BIOMASS, g m <sup>-2</sup>	
		AP		2-WK PE		4-WK PE		8-WK PE	
		COT	SOY	COT	SOY	COT	SOY	COT	SOY
NT-ROLL NO HERB*	NO PRE	2.9	7.5	17.4	14.8	29.6	29.4	53.6	54.7
SST-ROLL NO HERB**	NO PRE	3.0	--	17.4	--	27.0	--	120.4	--
NT-ROLL+GLY <sup>§</sup> ONLY	NO PRE	1.2	0.5	10.5	23.4	22.7	34.2	147.6	135.9
SST-ROLL+GLY ONLY	NO PRE	1.1	--	16.8	--	23.4	--	115.9	--
NT-ROLL+GLY+PRE HERB	+PRE	0.1	0	0.1	0.7	0.4	1.5	5.9	3.2
SST-ROLL+GLY+PRE HERB	+PRE	0	--	0.3	--	0.8	--	3.5	--
NT-ROLL+GLY+PRE+POST HERB	+PRE	0	0	0	0.7	1.0	0.1	1.3	0
SST-ROLL+GLY+PRE+POST HERB	+PRE	0.3	--	1.3	--	0.6	--	3.2	--
NT-NO ROLL+GLY+PRE+POST HERB	+PRE	0	0	0.1	1.0	0.2	1.3	0.1	0.3
CDT-GLY+PRE+POST HERB***	+PRE	0	0	0.9	1.8	0.2	3.3	0.3	7.2

<sup>‡</sup>Contrast of six treatments that received a pre- and/or post-emergence residual herbicide vs. four treatments without herbicide.

\* NT=no-tillage; \*\*SST= subsoil strip-tillage; \*\*\*CDT=clean disk tillage; § GLY=Glyphosate

## Results

### Weeds

Overall, the rye cover crop effectively suppressed growth of winter annual weeds. However, some cool-season brassicas became established in the rye and consequently, weed counts at planting tended to be higher in terms of the number of weeds surviving the rolling operation, when a burndown herbicide was not applied before rolling (**Table 1**). This produced significant ( $p < .0001$ ) differences in weed

(*Ipomoea* spp.), lambs quarters (*Chenopodium album*), eclipta (*Eclipta alba*), broadleaf signal grass (*Brachiaria platyphylla*), and carpetweed (*Mollugo verticillata*). The pressed, high-density residue also posed little challenge to seedling perennials such as pokeweed (*Phytolacca americana*), and curly dock (*Rumex crispus*). Short-lived perennials like horseweed (*Conyza canadensis*) and dog fennel (*Eupatorium capillofolium*) also were unfazed by the rolling strategy.

[3]

**Rye Residue**

Rye biomass yields are summarized in **Table 2**. In all cases, decomposition of the rye residue was fitted to a single exponential model. A significantly greater ( $p=0.0001$ ) rate of decomposition was observed in 2004 compared to 2005 (**Figure 1**). The year-to-year difference in decomposition rate can probably be explained by a difference in seasonal rainfall.


**Table 2.** Rye biomass yield ( $Mg\ ha^{-1}$ ) for two cropping seasons and two crop studies.

YEAR	CROP	MEAN	RANGE
2004	COT	4.97	4.50-5.76
2004	SOY	7.10	6.76-8.92
2005	COT	6.56	4.75-8.51
2005	SOY	6.54	4.95-8.47



**Figure 2.** Broadleaf signal grass emerging from a rye cover in the crop interrow. This plot received a glyphosate treatment at burndown, but no pre-emergence herbicide. Photo taken by this author 13 days after planting.

In 2004 nearly 30 inches of rain were recorded at the research station between May-September, whereas in 2005, 22 inches of rain fell during the same period. Model predictions indicated that four weeks AP approximately 67 and 78 per cent of the original residue remained on the soil surface in 2004 and 2005 respectively. In each crop and cropping season the quantity of residue produced was enough to provide 100% soil cover. Never the less, weeds began germinating <2 weeks after planting, and weed density increased progressively over the observation period (**Figure 2**). This suggests that the weed suppressive qualities of a rye mulch are ephemeral, even when high densities are achieved<sup>1</sup>.



**Figure 1.** Rye residue decomposition over 16 weeks (cotton trial data shown).

**Soil Moisture**

Volumetric soil moisture content differed as a function of both treatment row position ( $P=0.03$ ) and depth ( $p=0.001$ ). In general, differences in soil moisture in the surface 15 cm


were not significant across treatments and row positions. Comparison of row positions in subsoil strip-tillage showed greater soil moisture depletion in the tilled strip vs. the undisturbed interrow. Of greater consequence was the difference in soil moisture content between treatment row position with increasing depth (**Figures 3 and 4**). In plots where herbicide was not applied, and the cover crop terminated with the mechanical roller-crimper, soil moisture was significantly depleted in the no-tillage and strip-tillage interrow position at  $\geq 30$  cm deep, compared to where mechanical rolling was combined with a burndown herbicide to terminate cover crop growth. To express this contrast in soil moisture depletion in more practical terms:  $0.185\ cm^3\ H_2O$  per  $cm^3$  of soil (mean interrow moisture content of the ROLL NO HERB NT-COTTON treatment, 30 to 45-cm depth; see **Figure 3** pointer), is equivalent to 30,160 gallons  $H_2O$  per acre furrow slice (AFS). On the other hand, where the cover crop was chemically killed, mean moisture content in the 30 to 45-cm depth was  $0.252\ cm^3\ H_2O$  per  $cm^3$  of soil (mean interrow moisture content of the ROLL+GLY+PRE+POST NT-COTTON treatment; see **Figure 3** pointer), or 41,080 gallons per AFS. The difference is 10,920 gallons per AFS, or 0.40 inches of water per acre (1 acre inch = 27,154 gallons). The significance of this depletion is obvious in terms of available moisture going into the main cropping season. Since the crimping blades on the roller-crimper suppressed, but did not actually kill the cover crop immediately, it stands to reason that moisture uptake by the cover crop would still be occurring after rolling. And, since more winter annuals survived the roll/no-herbicide residue management strategy, this would tend to amplify the moisture depletion problem.

Here, an interesting question arises: Why did the mechanically killed rye preferentially extract moisture from the deeper soil zones, rather than near the surface where


rev: 05 Dec 2012

<sup>1</sup>Rye (and other crop) residues contain chemicals that, when extracted and assayed in the laboratory, have been shown to inhibit weed germination via 'allelopathy'. However, when these chemicals are leached into the soil they are likely diluted and undergo rapid microbial decomposition. Hence it is difficult to separate physical vs. chemical effects when assessing weed suppression by cover crops.

[4]



**Figure 3.** Above, tillage and method of residue management effects on early-season soil moisture in cotton (mean ± standard error).



**Figure 4.** Tillage and method of residue management effects on early-season soil moisture in soybean (mean ± standard error).

the mass of small grain roots are primarily concentrated? The answer isn't clear from this data. Some evaporation from the soil surface may have been occurring, causing soil moisture to equalize more rapidly in the prime root zone. The two sites had texturally dissimilar subsoil (loamy in soybeans and sandy in cotton) as noted in excavating the soil cores. But, in terms of soil moisture depletion, the response to non-chemical, mechanical growth termination in cotton and soybean plots was similar and may point to a potential drawback of the mechanical roller-crimper in organic cropping systems.

*Yield*

The effect of different weed management programs on yield was highly significant ( $p < .0001$ ) in both years. Our interpretation of the data suggests that high-density, surface pressed residue coupled with no-tillage, did not constitute an effective barrier against early-season weeds equal to that of a pre-emergent residual herbicide. In general, as herbicide input was sequentially reduced, yields declined lock-step in both crops (**Table 3**). When pre- and post-emergence herbicides were included as part of the weed management program, yields of cotton lint and soybean were not significantly different between tillage system and method of residue management (roll vs. no roll). Yield response to subsoiling in cotton was inconsistent; a modest gain (+247 kg ha<sup>-1</sup> lint) over no-till was noted in 2004 but none in 2005. The lack of response to sub-soiling in 2005 may have been due to early-season infection in some strip-tillage plots by *Phythium* root rot. Also, moisture was not limiting during most of the growing season in 2004, whereas in 2005 moisture was limiting only in the latter part of the growing season, so overall the effect of sub-soiling may have been attenuated.

**Table 3.** Cotton lint and soybean yield response to tillage and method of weed management over two cropping seasons.

TREATMENT	CONTRAST‡	COTTON LINT YIELD Kg ha <sup>-1</sup> (Lb /Acre)	SOYBEAN YIELD Mg ha <sup>-1</sup> (Bu/Acre)
NT- ROLL NO HERB*	NO PRE	327 (292)	2.15 (32)
SST- ROLL NO HERB**	NO PRE	321 (287)	--
NT- ROLL+GLY§ ONLY	NO PRE	396 (354)	2.22 (33)
SST- ROLL+GLY ONLY	NO PRE	520 (464)	--
NT- ROLL+GLY+PRE HERB	+PRE	1014 (905)	2.49 (37)
SST- ROLL+GLY+PRE HERB	+PRE	1196 (1068)	--
NT- ROLL+GLY+PRE+POST HERB	+PRE	1270 (1134)	2.82 (42)
SST- ROLL+GLY+PRE+POST HERB	+PRE	1334 (1191)	--
NT- NO ROLL+GLY+PRE+POST HERB	+PRE	1250 (1116)	3.09 (46)
CDT- GLY+PRE+POST HERB***	+PRE	1300 (1161)	2.96 (44)

‡Contrast of six treatments that received a pre- and/or post-emergence herbicide vs. four treatments without herbicide.

\*NT= no-tillage; \*\*SST=subsoil strip tillage; \*\*\* CDT=Clean disk tillage; § GLY=Glyphosate

Perhaps the most surprising outcome was the success we achieved planting into standing rye treated with glyphosate (Figures 5 and 6). Despite some initial reduction in stand, soybean and cotton lint yield was equal to, or better than, the yield obtained by mechanical rolling in both years.



**Figure 5.** Above, planting soybean no-till into standing rye residue. At right, **Figure 6** shows soybean growing in a dense thicket of dry, standing residue. Bean plants were a bit 'leggy' in the juvenile stage but matured normally.



### What Did We Learn?

Our research group's primary objective is evaluating different approaches to reducing or eliminating tillage, while creating a plant rooting environment in which crops can express their maximum yield and quality potential. Integrating tillage, residue management and pest control programs in ways that are flexible and mutually sustaining are important objectives. Following are the key points we realized from this project:

- Row crops can be successfully grown under high-residue conservation tillage using the mechanical roller-crimper.
- High-density residue and residue orientation were not as effective as herbicide in suppressing annual or seedling perennial weeds in a no-tillage or subsoil strip-tillage planting system.
- When herbicide input was reduced, there was a yield penalty regardless of tillage system.
- Mechanical cover crop rolling without glyphosate application was less effective in terminating rye growth as a glyphosate application followed by cover crop rolling.
- Mechanical rolling of living cover crops may impose soil moisture constraints on the principal crop, depending on effectiveness of the kill.
- Rolling and crimping a rye cover crop is not necessary to achieve good production in high residue, no-tillage soybean or cotton planting systems.

### Looking Forward

Soil degradation is a problem that continues to stifle agricultural productivity throughout the southeastern U.S. Piedmont and Coastal Plain regions. Much of the problem can be traced to exhaustive tillage, lack of adequate residue cover, and uncontrolled traffic in the field. Intensifying cropping systems through the use of winter annual cover crops is desirable as means of enhancing soil organic matter and realizing improvements in the plant rooting environment. Other factors such as narrow row spacing also may aid in limiting opportunities for weeds by increasing competition for light, moisture, and nutrients. Future work should therefore focus on continued integration of tillage, residue management, and weed control with attention to the following points:

- Field trials of a longer duration than two years are needed to evaluate the establishment effects of high residue systems. Providing that timely weed control is implemented for a period during the establishment phase of the cropping system, continuous no-tillage or minimum tillage coupled with narrow row spacing and full residue cover prior to canopy closure may ultimately mature into a production system that relies on fewer herbicides for weed management.

- Convenience and time-saving characteristics will continue to be a major concern for producers when it comes to cover crop residue management. The present two- and four-row mechanical roller-crimpers may be adequate for small producers; however, in order to expand adoption of the mechanical roller-crimper it will be necessary to alter its basic design to accommodate larger (eight to twelve+ row) planters as well as minimize vibration felt by the tractor operator. The present single or tandem drum arrangement of rollers cannot do this without becoming excessively massive and unwieldy. Design considerations should focus on smaller, in-row rollers that may be attached directly to the planter unit in a way similar to row cleaners and other residue management tools. In addition, if cover crop herbicides such as glyphosate are to be reduced or eliminated, the roller-crimper design must optimize termination of growth.
- Overall the trend in agriculture worldwide is driving toward increased productivity through increased cropping intensity per unit of land area. High residue conservation tillage is likely to impact the balance of nutrients and available soil moisture in the near term, as well as over multiple cropping seasons. Future work should therefore be directed toward long term assessment of profile moisture and nutrients in different cropping systems in the prime rooting zone as well as in the subsoil.
- The verdict is still out over the inherent benefits associated with rolling cover crops. Our data showed that mean soybean grain and cotton lint yield over two seasons did not differ significantly in response to the method of residue management. There may however be other merits and/or drawbacks associated with mechanical rolling that we did not perceive. Thus, we conclude that future work to evaluate the mechanical roller is justified.

---

### Acknowledgements

This project would not have been possible without the unstinted support of the farm crew at the Center for Environmental Farming Systems, Goldsboro, NC. Thanks also to Michael G. Waggoner, Department of Soil Science, N.C. State University and Keith R. Baldwin, N.C. A&T University, for their generous financial support of this project.

Prepared by:

Robert Walters  
 Department of Soil Science  
 North Carolina State University  
 CB 7619  
 Raleigh, NC 27695  
 Email: [robert\\_walters@ncsu.edu](mailto:robert_walters@ncsu.edu)